CONSIDERATIONS

ON

BILIOUS DISEASES:

AND

OF THE LIVER, AND THE GALL BLADDER.

THE SECOND EDITION,

By JOHN ANDREE, M.D.

" AD UTILITATEN VITA, ONNIA COMILIA PACTAQUE

LONDON:

Printed for J. MURRAY, No. 32, Fleet-Street; J. Johnson, No. 72, St. Pani's-Church-Yard; W. CLARKE, No. 38, New Bond-Street; P. HILL, and WATSON, ELDER and Co. Edinburgh; A. CLARKE and T. SIMSON, Hertford.

MPCCLERES.

COMSIDERATIONS

n u

BILLIOUS DISEASES.

AND

SOME CARLICLAR ASPROLATING OF THE LAVER, AND ER.

THE CARROND EDITION



"AD LITTER STREET ON TA CONSIGNATION OF THE CONTROL OF THE CONTROL

CAN O O WOLL

Applied for themselve, N. co. Flooration I.

foreigner, NE co. Northanie Complex Yante W. Charles,

vo. co. Nord Senere. P. Herrison and Warrawa,
i. nord so. Conservation of Conservation Servation.

. * *** *****

1 10 -7

so Falling or the equility knowledge of

the kind of difeates, in the common

P. RuinE of Fra A Come

anthing the peers affected by differing

or many who had been adheded with such

disaders, who should distill that

forme of the epiphon contained in the

following nurse will tend to the roller

BILIOUS Diseases are among the most common of the chronic distempers of the inhabitants of England, and therefore merit the peculiar attention of medical practitioners. During twelve years, I had too much reason to lament the imperfect state of this part of medical knowledge, having been nearly so long subject to a bilious disorder, undescribed by any authors, and opprobrious to my own, and learned friends kind medical aid.

Failing

Failing in the requisite knowledge of these kind of diseases, in the common course of medical learning, my next recourse was to anatomical investigation; by embracing every opportunity of eximining the parts affected by diffection of many who had been afflicted with fuch disorders .- Having flattered myself that some of the opinions contained in the following pages will tend to the relief of mankind in such diseases, I folt it p duty incumbent on me to submit them to public inspection in this, still far from perfect fate, fatisfying myfelf with " mens fibi conscia recti." resoidibary las

I had too much reason to lanent the The Dropfy being not unfrequently preceded by bilious disease, I trust the introduction of the following remark will not be judged wholly irrelevant; to wit, that when it has continued for a certain length of time, that is to fay, Failing

is become a chronic distemper, "the hasty evacuation of the water," whether by medicine or surgical operation, "will precipitate the death of the patient; apparently from inanition."—Also that the digitalis, from it's baneful effects," is to be classed among the deleterious medicines, and therefore should not be used until the common diuretics have been administered; among which the dried squill claims the pre-eminence.

Hertford, May, 1790.

[•] See a paper on this subject by Dr. Lettsom, in Memoirs of the Medical Society of London, Vol. II.

infly evacuation of the water," whather in five condition of the place, "whather in condition of the place, and the proception of the place of the place, and the proception of the condition, and the condition of the condition o

ERRATA

Page 35. hall ling, for Cullen's read Cullen,
Page 41. last line but one, for his read sire

tale alteriorated at 100 food halfs to expense on a

A fire technology that the telephone when

lower part of the ribs and their cartiar ginous appendages. It covers the upper part of the flourachy and formed the displacement of the bowels, it adheres to the displacement of the bowest expanded muscle, separating the cavity of dibe twoisid for notiniqo starating attantide the cavity of the cast that the cavity of the cast that the part of the cavity of the cast that the part of the cast that the cavity of the cast that the part of the cavity of the cast that the part of the cast that the cavity and the cast that the cavity and the cavity and the cavity and the cavity and the cavity beautiful the cavity of the cavity and the cavity beautiful the cavity and the cavity and the cavity of the cavity and the cavity of the

ender these general heads it shall be my ender vour to elucidate this subject. Previsous to which it may not be improper to give a brief anatomical description of the liver, the gall bladder and ductare soul and such an

body; of a dusky red color. It is fituated

IIA B princi-

principally on the right fide, and upper part of the abdomen, being covered by the lower part of the ribs and their cartilaginous appendages. It covers the upper part of the stomach, and some of the bowels; it adheres to the diaphragm, a thin expanded muscle, separating the cavity of the cheft and the lungs, from the abdomen. It is tied to the breast bone by a broad ligament, and to the navel by a round one: It is nourished by the arteria hepatica: but its blood vessels, from which the bile is fecreted, are the branches of the vena portarum, which distributes its blood like an artery, in order for the fecretion of the bile. Its weight in a middle fized man is about three pounds twelve ounces tall out

bag, feated in the hollow part of the liver; ending in a duct called cyfticus, which is united to another duct coming from the liver, named hepaticus, land from thefe, thus joined, commences the ductus communis cholidochus; which enters the first gut near the stomach. The shall a to the ductus and from the liver.

All the bile secreted in the liver passes through the hepatic duct in its course to the gut, and a part regurgitates into the gall bladder. and a bladder and bladder.

The first subject is a redundancy of bile.

This has ever been considered as one cause of diarrhea's, more common in warm climates than in England, where I think it is frequently productive of a confitutional lax habit of body.

Another effect of a redundancy of bile is a vomiting of that fluid, appearing as the effort of the constitution to free itself of the redundant quantity of bile. Of this I have seen some, besides the ensuing instances, ter-

sile. This continued for three years, with-

B 2

W. A.

minating

I opened the abdomen of a healthy dog, tied up the biliary ducts, then sewed up the wound in the belly, and the animal recovered. Ten days afterwards the dog was killed, and I found the gall bladder perfectly flaccid and empty.—This experiment shews, that all the bile found in the gall bladder gets into that vessel by regurgitation, through the biliary ducts; and that there are no canals leading immediately from the liver to it, as supposed by some learned authors.

-middlibiration de deposite duck in its courtered the guit, and a part regurgitates into the

This discharge of bile becomes in some instances habitual; I have also remarked it to become periodical; and most instally to happen either on a full, or on an empty stomach; for example, at first riling out of bed, or after eating a full meal; instances of each of which I have known. This symptom may continue many months; perhaps years, without materially injuring the constitution.

J. H. Esq; aged twenty-five years, of a healthy and strong constitution, in 1773, began daily to vomit soon after dinner much of what he had cat, and some clear bile. This continued for three years, without any evident injury to his health, and then disappeared. His father is gouty, and himself has had several attacks of disorders attended with plethora. This symptom appeared in this case as the effort of the constitution, to throw off a surplus of bile.

W. A.

differentia

W. A. S. Elq. aged forty, has palfed at different times fix years in India, where he had repeated attacks of bilious difeafe. He enjoys, when living very cautiously in both cating and drinking, good health. Fat, and butter, taken freely, or a hearty meal of rich high fauced victuals, always produce a bilious vomiting. The remedy this gentleman uses is an aloctic purge, whenever his bowels are not in a rather lax state. For costiveness in him, brings on oppression at the stomach, head-ach, and some appearance of jaundice, especially about the eyes.

L. W. an unmarried lady aged twenty, was feized in 1785, with vomiting of bile in the morning, at first rising in the bed, which returned daily for five months. She then had frequent attacks of the same in the afternoon. This having continued near a twelvemonth ceased, costiveness ensued, and a dropsy put a fatal period to the disease.—In this case it is highly probable these ill consequences might have been prevented, had a proper course of medianes.

PORKETS

cines been adopted when the bilious discharge scased, which should have been joined with or followed by tonics.

er over when firms were cautiouly in

Another effect arising from this cause is a reassumption of bile into the circulation, demonstrated by its producing symptoms of jaundice recurring frequently on the commitment of any material error in diet. This is shewn by its producing yellowness of the eyes, and skin, particularly about the temples; from a continuance of such cause, a confirmed jaundice may also ensue.

Mr. N. Silversmith in the Strand, aged about thirty-five, in August 1786, consulted me, he had then been four months afflicted with a confirmed jaundice; for which, according to his own expression, he had taken his hat full of medicines without receiving any benefit. His health was otherwise good; he was rather costive, but had very little uncasiness, nor had he everseltany acute pains or spasms in the right side or near the liver, so as to give any rea-

fon for supposing there were biliary cal-

And I am Arongly perfugied that I bak

After maturely considering this case, it occurred to me that the disorder might be caused by an excess of bile. I therefore prescribed for him, pills of aloe, sapo, rhabarb. & gum. ammoniac. in order to keep the bowels gently lax. And thrice in the day to take a large tea spoonful of elix. vitr. acid. and tinctur. stomach. each equal parts. Also to leave off butter and fat, and to eat plentifully of china oranges, and other mild acid fruit.

He found benefit from the medicines in a few days: and his jaundice was completely gone in a fortnight.

on experiments, has been toned never to

Laddle proditioners is the Well ladies

He then left off the medicines, and his disorder returned in September, when it was again cured by the same means. The beginning of December he had another relapse, and was then cured by the acid drops alone, and regetable acids.

Thus

B. 4 want to B. 4 want to Si obi VA re-

A redundancy of bile seems also to be capable of inducing muscular debility.— And I am strongly persuaded that some obstinate eruptions on the skin arise from this cause; some instances of which I have seen.—An excess of bile (when become acrid or putrid) and absorbed into the blood, is also regarded as the exciting cause of bilious severs, not uncommon in England in spring, autumn, and hot and dry summers, and one of the most common diseases of Europeans in the East and West Indies.

English practitioners in the West Indies talk of an acid and an alcaline bile, alledging that in the former, acids encrease the disorders, in the latter are serviceable. Bile, on experiments, has been found never to become acid, nor does it become putrid in less time than the other animal secretions. The acid ejected by vomit in such cases, must therefore have been secreted in the stomach, or formed by the aliment previously contained therein, or in its vicinity.

That

Vide Boerhaave's Academical Lectures.

That acids disagree with some disorders of the bilious kind is certain; but when it is confidered that the stomach does in fome cases actually secrete a superabundant acid, it will tend strongly towards the overthrowing of this theory.-The practice, however, is judicious, being to procure an immediate and effectual discharge of the offending fluids by tartar emetic: and then to throw in the bark boldly, which they alledge is found by experience to be the only fafe method of cure, and prevention of bilious fevers of the most fatal kind. That the bile does fometimes become of an irritating quality, so as to produce diarrhæa and some fevers, I think highly probable; though in my opinion less frequently so than is imagined. By much the greater number, I suppose, arise from the retention of a quantity of indigested food, which becoming highly rancid or putrid, is the true cause of such fevers. For fince it is certain that the bile out of the human body is very little prone to become putrid, it will, I think, on a little reflection, be evident, that animal substances, taken into the stomach, are much more likely to become putrid therein, than the bile. In the cure of such fevers the physician's attention should ever be fixed on the cause of the disease. The present practice of physic in these disorders being judicious, it will not be for me to add much on that subject.

In addition, however, to the ordinary remedies, I would recommend the liberal use of acids duly diluted, that is, for the patients common beverage, on the principle of their partially decompounding the bile, and thereby lessening the symptoms. The human bile is a sluid of an oleaginous viscidity, of the nature of a liquid animal sope, it is consequently of an alkaline nature, which the acids have a tendency to decompound, and thereby render inactive: the good effects of the above regimen I experienced in my own case. The liberal use of mild acids is also found of the greatest benefit in such diseases in the West

Indies.

Bilis sapo est. Haller, primæ lineæ physiol.

Indies, where it is even customary for patients to drink sherbet, in order to promote the operation of cathartic medicines.

In the East Indies also, mild acids have been found useful in bilious diseases. In Dr. G. Fordyce's elements of physic are these words: a moderate use of sour fruits, in warm summers and hot climates, tends also to prevent the diarrhæa, dysentery, or cholera morbus,

But should acids disagree, which may arise from a superabundance of the acid secreted in the stomach, plentiful dilution with water is to be recommended, from its readily uniting with the bile, which would be thereby rendered less active or pungent.

Imperial water, old hock and water, cyder, wine duly diluted, and whey or tea may be allowed. Malt liquor usually does harm, and spirits should be strictly forbidden.

It is also to be considered that moderation in the quantity of acid taken is necesfary, for if more is taken in than what is necessary for the decomposition of the bile and other juices, the acidity of the fuperfluous quantity remains; which might produce the ordinary symptoms of acidity in the stomach and bowels. Whereas, on the contrary, if the intestines are overcharged with animal food, it will tend much to encrease the symptoms in this kind of bilious habit, either by producing a greater quantity of bile, or of a bad quality, and having a confiderably stronger tendency to become tenacious and putrescent, than it in a healthy state should be indued with driv online vibes and

In support of this doctrine, let it be observed that Europeans, living in hot countries, who eat much of animal food, are by far more subject to bilious diseases than the natives, whose diet consists principally of vegetables and grain, and whose drink is water.

would be thereby rene

Moderation in the quantity then is indicated, and animal food should be allowed only once in every day, and probably one or two days abstinence in every week from such diet would be of surther advantage. The remainder of the food should be attenuant, (nutritive, farinaceous diet, port wine and madeira are also strongly recommended,*) easy of digestion, and mostly vegetable: whereas salted meats, most kinds of sish, sweet cakes, butter, especially when melted, and fat of meat, are hurtful.

But another most essential circumstance for such subjects to attend to, is the due and constant preservation of a moderately lax habit of body. If this can be accomplished by diet, it is to be preserved to medicine. But general rules for this can not be laid down, because every such partient, duely attentive to his own health, may from observation be enabled to ascertain the kind of diet most conducive

It however mould be adapted to the

one case which did well, the patient was led by tade to

[.] Dr. Blane on Diseases of Seamen.

to this end. Substituting honey (which, where it agrees, is one of the most falutary ingredients in diet) instead of butter, I have known answer this purpose well; and the drinking some warm water gruel at going to bed has tended much to this; the total abstinence from butter and fat should also be tried.

Exercise is another, and a most powerful healthy stimulus to the liver, bowels, and digestive organs.

Gronely recommended.*) esto of direct

It however should be adapted to the patient's strength and its effects.—Riding

of meate are hurtfull.

• A friend of mine told me, that in September, 1787, being in Norfolk on a shooting party, he was attacked with a cholera morbus, the vomiting continued at intervals during the first day, and then ceased. On the ensuing two days he had a bilious slux, having, he thinks, not less than thirty dejections on the first day, and twenty the second. He lived upon thin oatmeal gruel, which he found to agree better with him than any other diet: and the disease went off without the aid of medicine.

In Dr. Blane's Observations on the Diseases of Seamen, are these words on the subject yellow sever; "In one case which did well, the patient was led by taste to prefer warm water gruel to every thing else, and the great quantity he drank," &c. on horseback is with justice accounted the best exercise for the bilious; this should also be joined with change of air; that of a dry and bracing kind is to be preferred.

I have known attracks of bilious difeate

Any exercise is adviseable which will necessarily induce the patient frequently to bend the body in various directions *. Swinging, (lately recommended for confumptions) might be tried with some prospect of advantage, on the same principle; as an exercise well adapted to affist such cases in a harmless manner. Whereas I have known in myself violent jolting of the body, for example on a hard trotting horse, produce pain, and exaggerate bilious fymptoms: and if there were rugged biliary calculi in the gall bladder or ducts, might even move them fo as to irritate and bring on violent spasmodic pains in the parts affected, and even be attended

flered.

without flimulating, are to be admini-

Of fuch medicinal use has muscular exercise been considered, that Fuller and others have written on medicinal gymnastics.

with fome danger of lacerating the

also be joined with change of air; that

It is also certain that the mind has a powerful influence upon the biliary parts. I have known attacks of bilious disease repeatedly brought on by vexation, or anxiety of mind.

to bend the body in various directions

Hildanus has remarked that blows of the head, have brought on fatal diseases of the liver. A bilious vomiting is a symptom following often, very quickly after accidental injuries of the brain, and sun strokes in the East Indies; may not therefore the liver and its ducts, as well as the stomach be supposed to sympathize with the brain?

The bowels it has been faid should be kept open. If this cannot be accomplished by diet, such purgatives as will act without stimulating, are to be admini-

lovi in the call bladder or ducker

ftered

The wife of a gentleman of the faculty in London died of a rupture of the gall bladder, from laceration with a gall stone, in 1786.

stered. But no general rule can be laid down for this; various remedies agreeing with different persons. Castor oil is a valuable medicine for this purpose. Manna with falts warmed with some cordial water, is used with advantage by some, rhubarb possessing a tonic effect upon the bowels is to be recommended, and the aloetic kind for its certainty of operating is in some cases to be chosen: calomel. sope and rhubarb given together, are said tobring away more bile than any other purgative. But the best laxatives are purging mineral waters, of which the kindliest in their operation are those of Cheltenham, being indued with the peculiar property of keeping the body open without debilitating.

Sailing, in obstinate cases might also be recommended, as productive of a sickness attended with extreme languor, or atony equal to the action of opium or a deliquium from the loss of blood, without so much debilitating the system: much benefit

benefit may also be derived from the brace ing effects of the sea air.

The last indication of cure is tostrengthen the tone of the vessels of the liver. For this purpose chalybeates may be recommended; the Peruvian bark has been advised, but is found not to act kindly in bilious diseases; the columbo is now preferred, especially in India, where bilious diseases are more common than here. The quassia should also be tried, being very beneficial in some stomach diseases.

This bilious habit of body is also in fome instances constitutional and not productive of disease, attendant upon what is called the hot constitution, that is, vigorous and irritable mind, with a warm surface of the body, quick circulation, and tendency to instammatory diseases.

Another extraordinary bilious disease, is the enlarged gall bladder. Of this disorder I have the following case to relate, through favor of my friend Mr. Cline, surgeon to St. Thomas's Hospital.

S. M.

S. M. aged fixteen, had a troublesome cough for feveral months, attended with pain in the right hypochondriac region. On the same fide a tumor gradually formed, and diftended the cavity of the abdomen: a fluctuation being distinctly felt. a trochar was introduced. By this operation more than twenty ounces of bilious fluid was discharged. During the evacuation, he complained of great pain in the part and in his right shoulder: symptoms of inflammation foon followed, and he died on the seventh day after the opebladder cight pounds of thick biknoits

He had been in an ill state of health about twelve months preceding his death, but without any suspicion of obstruction to the passage of the bile; for his skin had not been discoloured nor his stools apparently altered inages to stol ; bodicalab ands

hard and clay coloured thools, at times of Dr. Cheston of Glocester, examined the body, and found the gall bladder, containing about two quarts of bile, extend-

ing from its usual situation down to the pelvis, and adhering to the peritonœum omentum and part of the stomach, all of which were inslamed. The biliary ducts were greatly inslamed, except where the ductus communis cholidochus enters the duodenum, which part was contracted, but admitted the bile, with some difficulty, to be pressed into the intestine.

Van Swieten, in his commentaries on Boerhaave's aphorisms, relates a case of a lad twelve years of age, in whose gall bladder eight pounds of thick bile were found.

cuation, become lagred of great pain in

The fecond subject of consideration is a diminished secretion of bile.

He had been in an ill flage of health

The symptoms of want of bile may be thus described; loss of appetite, costiveness, hard and clay coloured stools, at times of a putrid fætor, a peculiarly sickly paleness of the face, loss of slesh, indigestion, attended with slatulencies, a langour of the body and mind, and spasmodic asthma.

It is also productive of a peculiarly morbid complexion, neither yellow, confumptive, nor pallid, but of such a peculiar ill aspect, (a kind of leaden color,) as I suppose is not describable by words, but which an attentive observer might ever remember, having once thoroughly noticed.

I am strongly of opinion that many (perhaps it will be found that the greater number of) asthma's are caused by disease of the liver. In India, bilious disease is accounted as the cause of many coughs: and the late Mr. Paseley, an ingenious and sensible practitioner in India, remarked, that in all confirmed diseases of the lungs, the liver was affected.—He even supposed every chronic distemper to originate from disease of the liver.—Another remark of the same gentleman was, that in all liver diseases the urine was high coloured.

Such are the symptoms too often seen in this island as the consequences of drink-

C 3.

ing, particularly spirits, and in such cases usually the harbingers of fatal dropsy, or consumption.

Some of these symptoms are attendant upon the schirrus liver, and may perhaps be sufficient to mark that distemper.—It is the common practice to examine by the hands externally on the abdomen, when an induration of the liver is suspected; on the idea that it is always then enlarged. The following cases however evince that such examination is not to be relied on; for it is by these proved that such disease may occur with a decrease of its size, in which situation it would not be perceptible by an examination during life with the hands.

This observation I made in 1774, on having twice found the liver diseased and tending to schirrus, and at the same time shrunk within the margin of the ribs, so as not to have been perceptible by manual examination during life.

J. G. aged fifty, after great fatigue from travelling, was seized with pain in the chest, cough and spitting, attended with shortness of breathing, particularly after eating. A spitting of blood then ensued, which however ceased a month before death, during which time he was free from pain in the chest, but could not lay on his right side.

A sudden profuse bleeding from the lungs, suffocated this patient. On dissection some slight disease of the lungs was discovered. The liver was indurated, diminished in size, of a paler color than natural, throughout its whole sustance: and the gall bladder was very much thickened.

B. J. aged forty, had been for many years subject to pains in the right hypochondrium, which returned periodically, and were often accompanied with vomitings and asthma. Among other less morbid appearances, such as serum between the membranes of the brain and its

C 4 ventri-

ventricles: the liver was found hardened, but not enlarged.

Another kind of indurated and diminished liver which I have also met with, is that in which it's surface is raised in half rounded eminences, appearing like granulations, covered with the proper membrane of the liver.

In one of these cases, the particulars of which I have inadvertently lost, the patient died dropsical.

A gentleman aged fifty-nine, was attacked in May, 1787, with lassifude, loss of appetite, and pains in the loins extending towards sometimes the one, sometimes the other shoulder, the urine also was very thick and red, with a yellow tinge, and deposited a thick red sediment; it at times was charged with a ropey mucus, somewhat resembling matter. The pulse was small, and for the most part slow. These symptoms were regarded as what is vulgarly called a breaking up of

the constitution: and palliatives only were prescribed.

Having discovered a dingey yellowness of the skin and eyes, I advised that the case should be treated as bilious: and prescribed pills of aloes with gum ammoniac. soap and rhubarb.—After a lingering disease of five months, the unhappy patient's sufferings terminated in death. During this period he was often attacked with asthma. It was spasmodic: attacking him at various intervals, and admitted of no relief but from opiates.

During the progress of this disease, various medicinal plans were pursued, such as sedatives, balsamics, mercurials in small doses, cicuta, solvents of calculus, and laxatives: with change of air.

Another remarkable symptom in this case was the elevation of the shoulders, observed as a symptom of diseased liver by Bontius, whose translator, stiles it, prominence of the shoulders. Also by

Dr. Girdlestone, and others of the mo-

On examination after death when the abdomen was opened, the omentum, stomach and bowels, also the kidneys and ureters were free from disease, as were the other parts, excepting the liver, which was reduced to about one half it's natural size, and remarkably indurated; and at the outer part to the depth of three-eighths of an inch positively as hard as griftle, and the remainder completely schirrus.*

It

* Boerhaave in the body of an officer, who had gone through all the degrees of the yellow, green, and black jaundice, found the liver scarce equal to a hand's breadth, and about the same thickness, but as tough as leather. Boerhaave's Academical Lectures.

Riolan relates that he found in a body which he opened at Paris, the liver fo small as scarcely to equal the fize of a kidney.

Van Swieten writes that he has fometimes found the whole liver schirrus, and contracted, or greatly diminished in bulk, juiceless, and tough like leather.

Dr. C. H. Velse, in his inaugural thesis at Leyden, 1742, relates a case in which the liver was hard, rigid,

It appeared then that it would have been impossible to have discovered by manual examination the nature of the disease of the liver in these cases, since it was contracted above the margin of the ribs, consequently covered by bone, and beyond the reach of the hand in an examination. The enlarged liver is, however, a much more common disease: * In the following case the symptoms were of the same kind

of diminished bulk, and befet with solid rough tubercles, over its whole convex surface.

Du Verney found the liver fo dried in half its fubflance as not to exceed the thickness of his thumb.

The late Dr. Hunter in his lectures, related a case in which the liver was schrirrus and much shrunk or distinished.

In Matthews's observations on hepatic diseases, page 202, are these words; the liver was found remarkably small and hard in a patient who died of hepatic dysentery.

The celebrated Ruysh found the liver in two instances of dropsical subjects, indurated and not enlarged.

* It has been found encreased in bulk, so as to weigh fourteen pounds four ounces, and extending over great part of the abdomen: and often to extend much below the margin of the ribs. An enlarged and indurated liver is one of the most common chronic diseases of those of the abdomen.

with

with the former, although the liver was enlarged.

A. W. aged forty years, had been long afflicted with a large hard tumor on the right fide of the abdomen, painful to the touch; attended with thirst, slow fever, difficulty of breathing and vomiting.

On examination after death, the tumor was discovered to be the liver grown to a great bulk, it's right lobe extending almost to the lower part of the belly.

This lobe in particular was univerfally indurated, as was the remainder of the liver, though not so much so in it's inner part. The coats of the gall bladder were thickened, it's cavity small, and containing some black, thick, viscid bile.

The lungs had no other diseased appearance, excepting some dark colored spots.—I forbear inserting more cases of this kind, because such may be found in various authors, attended with many of the

the most distressing symptoms above recited.

It appears to me that the symptoms already described, likewise violent pain in the kidneys extending up to the shoulder, urine of a deep yellowish red color, and depositing a sediment seeming to be red particles of blood, loss of appetite, indigestion, and statulencies in the stomach and bowels, were chiefly caused by the want of bile; but that these symptoms were to have been in part attributed to the diseased state of the liver, is also allowed.

The cases above recited, it is true ended satally; but had their cause been known, another and powerful remedy should have been used, viz. a regular course of mercurial frictions—Most of the above symptoms in a less degree, are not uncommon, especially in those who have suffered liver disease in hot countries: sometimes happening from inflammation

flammation of that gland, having left it in an indurated state.*

In addition to the above, as a palliative, I would recommend animal bile inspiffated, (that of the ox has been given,) in the form of pills, in order to supply the want of the human bile. I have within the last year prescribed it in two cases with success. In one of these cases, until this medicine was taken, during many preceding weeks, the patient voided no stools without clysters, and reaped the advantage of natural dejections by pills formed of the bile of ox inspissated.

Boerhaave, as I am credibly informed, cured an obstinate jaundice in the child

of the breath, in inspirations is a symptom of the inflammatory state of the liver: and that a deep hollow sounding cough is a symptom of an abcess in that gland."

The term obstruction is still familiar in physic, for example an induration of the liver is stiled an obstruction, an induration and enlargement of the mesenteric glands is called an obstruction of them; it is used in conformity to the Boerhaavian theory of instammation: but is repugnant to the known laws of the animal economy.

of an English merchant, with bile of the pike fish inspissated.

This has been found a successful medicine for epilepsy,* accompanied with a bloated and pale countenance, indicative of some visceral disease, probably of the liver.

The method of cure then should be, first to attempt the resolution of the induration of this gland, by the ordinary deobstruent and attenuant medicines; of which the aloetic kind, assisted with rhubarb, sope, and alkaline salts, may be first tried: which in an incipient state of congestion of the liver, will in some cases prove essection.

Strong or cold purgatives are dangerous: the former by their irritating quality might induce inflammation: and the latter I have known bring on a jaundice, once in

^{*} Vide Dr. Duncan's medical commentaries for 1788, page 162: ox bile recommended for epilepsy, by the late Dr. Quarin.

myself; also in another case when prescribed by a physician of eminence.*

The next plan which I think merits a trial: but to which much time should not be given up, is the cicuta, the extract, (fuccus spissatus,) which should be given twice a day, encreasing the dose gradually until it produces some pain or dizziness in the head: tartar emetic in small doses may be joined with this medicine, in order to render it more active. Much depends on the goodness of this medicine, and I hold it as an excellent rule, "that when hemlock, either in the form of powder or extract. has no sensible effects when taken to twenty grains for a dose, the medicine may be supposed to be impersect; and that, if it is continued, another parcel of it should be employed." Cullen's Materia Medica, vol. 2. p. 264.

But

^{*} Each of these cases happened in hot weather.—A due regard to the correcting the ill quality of the bile, previous to it's hasty evacuation, is therefore worthy of attention, especially at such seasons.

But the most effectual remedy in such cases, is a course of mercurial inunctions,* fo conducted as to excite a moderate falivation, or at least some degree of foreness of the mouth.-The frictions were at their first introduction among the British, made on the region of the liver: but it having been found that the act of rubbing-in the ointment over this diseased gland was of prejudice, the common mode of rubbing it into the lower extremities was substituted: and it's success has justified the practice. Mercurial purgatives are also to be interposed occasionally: for these, assisted with camphor, opium, or fuch other medicines as the fymptoms shall require, may effect a cure in an early state of the disease.

D Henry

The late celebrated Dr. Cullen in his lectures, with his well known candor, and the liberality of a great mind faid, we have lately learnt from the gentlemen practitioners in the East Indies, that the liberal use of mercurials is the only effectual means of discussing congestions in the liver. -It is also afferted that to the full proportion of nineteen in twenty of such cases are thus curable. Vide Matthews citat.

Henry Walker, aged thirty-three, a thin, healthy man, employed at the leather mills near Hertford, was attacked July 2, 1780, with a cold fit terminating in fever, but not accompanied with fuch privation of strength as occurs in the fimple continued fever. At the end of ten days the fever continued, and his eyes, forehead and temples were yellowish;on manual examination fome degree of tenderness to the touch and induration were perceived on the liver. A course of mercurial inunctions was immediately began (the pulse then exceeded 100) asfifted with calomel internally; the fymptoms gradually fubfided under this course, and the man was effectually cured in three weeks and a few days.

The practice of giving calomel in large doses in dropsies arising from diseased liver, was written of by Riverius above a century ago, as not uncommon then: he gives many cases which he cured with that practice. A few words may be necessary on the subject of mercurials, in order

order to obviate the prejudices conceived against them.

During full twenty years practice, and so long constant attention to the action of mercurials, I never have feen any fuch effects, as to induce me to believe it broke down the texture of the blood, or otherwife injured the constitution, when judiciously administered. In myself, when much debilitated by jaundice, in 1786, owing to congestion of the liver, and inert, probably viscid state of the bile, I took from three to five grains of calomel five nights in a week, for three weeks. which did not impair my strength .- Nor have I any reason to suppose that mercurials injure the folids. It is known from constant observation to stimulate various fecretory organs; as the falivary glands. exciting falivation; skin producing perspiration; intestines, diarrhæa; kidneys, urine; "and I am of opinion that we may add that it stimulates the liver, and encreases the secretion of bile."*

D 2 Solvens

^{*} Vide Cullen's Citat. article Hydrargyrus.

Solvens mineralis, a fublimation of arfenic, prepared at apothecaries hall, and highly extolled by a physician of eminence for it's good effects in schirri, might also be tried. It is to be given in doses of the tenth of a grain, twice a day, encreasing the dose gradually.

The use of the bath waters at Bath, is also a remedy highly deserving attention; where the circumstances of the case will admit of their use.

The medicinal treatment having been briefly spoken of, it may be proper now to treat on the diet best adapted for these cases.

The food should be nutrient and easy of digestion; of this kind, of slesh, are the (feræ natura, game) poultry, and such kind of sish as have the same qualities: of common food the choice will depend on the patient's experience of that which digests best, and sets easiest on the stomach. Salted meats, fat, butter, and lard, particularly

ticularly as introduced in pastry, and all condiments, should be avoided. However. fuch patients having feldom inclination or ability to eat much, even enough to preferve a due degree of strength in the constitution for it to combat the disease; the principal rule to be observed in their diet, is not to overcharge the stomach with a large quantity of food at any one time. The food should be attenuant: the drink may be whey, water, tea, barley water, occasionally taking some such cordial as wine, (with egg or spice when requisite) cyder may be advised, as may a light infusion of ginger, orange or lemon peel, camomile or gentian.

In many instances, bilious diseases having been brought on by gross feeding, and want of exercise, the article of diet in such, becomes a matter of the most serious consequence.—It is a very just observation that diet is capable of producing greater alteration in the human body than medicine, and with less injury to the constitution.

D₃ The

The late Dr. Huxham in the preface to his essay on fevers, says, on the subject of diet, " what we use by ounces and pounds cannot but confiderably affect us." And adds, that "Hippocrates and the ancients were very careful in prescribing a very exact regimen." In short, if error in diet is the cause, remove the cause and the effect ceases. I wish not to be misunderstood, as meaning that an indurated liver can be cured by any change of diet, but that by a proper diet its fymptoms may be alleviated. For the same author adds from Hippocrates (as his apology for not giving many formulæ of medicines,) " he that knows a disease, knows what is proper to cure it."

During either of the above courses, painful symptoms may be safely palliated even by opiates, provided the bowels are kept open.

Bitters it has been faid should be given, (tho' their safety when long continued is not thoroughly established:) as a stimulus

to the stomach they may tend to excite appetite, but from their tendency to excite contraction of the vessels, their action upon the biliary passages might tend to encrease the obstruction to the bile's free passage. But provided the disease should happily yield to medicine, towards the conclusion, the cautious use of bitters* with chalybeates may be with propriety for a time prescribed, in order to strengthen the weakened vessels of the liver. and digestive organs, and thereby prevent a relapse. Of chalybeates those mineral waters which are laxative, fuch are those of Tunbridge and Islington, are to be preferred to the medicinal preparations of feel. And the columbo root as a tonic and stomachic should be administered. It is afferted by practitioners of physic in India, that it will effect this purpose fully and with fafety: whereas the bark is by them reckoned productive of future obstructions.

^{*} There is some reason to suppose they are injurious to the eyes; and their long continued use certainly tends to weaken the stomach.

I cannot conclude this article without subjoining, that spirits should be religiously refrained from. The familiar experiment of soaking a piece of slesh in spirits, demonstrates their power of hardening animal slesh, and this effect in some degree must be allowed to take place upon the stomach and liver, in the living body.

The third subject for discussion is misplaced bile.

The morbus regius, aurigo, icterus, morbus arquatus, or jaundice, is the most obvious example of misplaced bile.

It consists of a mixture of the bile in the circulating mass of blood, distinguished by a yellow colour of the skin and eyes, attended with high coloured urine, staining linen yellow: costiveness and white hard stools, sometimes settid. It is usually productive of langour of body and mind, coldness or chilliness, loss

The pulse has been found remarkably flow in some instances of jaundice. The late Dr. Hunter found it in

loss of appetite, uneafiness in the right hypochondrium particularly on pressure, heat about the nostrils, and itching of the skin. In the advanced state every thing tafted feems bitter. The faliva and blood are tinged; and the whole contents of the abdomen and thorax. likewise the brain and the bones, have been found yellow on diffection. The eyes also as the poet remarks, "lurida præterea cernunt, quæcunque spectantur arquati," are sometimes so affected, that all objects appear yellow. This however was not fo in either of my three attacks of jaundice, nor have I ever met with an instance of it.+

one case only thirty-seven in a minute. I have now a dropsical case under my care, in which the pulse has been at thirty-three in a minute, full and regular, the man quite sensible, and is nearly cured.

From these circumstances may we not allow, that one use of the bile is, to affift in generating heat in the body?

† This opinion was maintained by Galen and others of the ancients. And two cases in which it occurred are mentioned by Hoffman, one by Sardona in his Aphorisms de cognoscend. et cur. morb. and two by Dr. Percival, in the second vol. of his Memoirs of the Med. Society of London.

This

This disease originates from various causes, the most obvious of which are mechanical obstructions of the biliary ducts. These may arise from biliary calculi; from obstructions in the biliary follicles of the liver; from induration or schirrus of some adjacent part pressing on the ducts: and another, and I suspect more common cause, to be a contraction of the gall ducts themselves: viscidity of the bile may also be admitted as another cause.

That the gall bladder and its ducts are liable to generate concretions called calculi, has been known from before the time of Galen. The calculus felleus is a more frequent disease than is imagined. These stones are of all varieties of figures and divers colours; they are however usually of a triangular shape when many in number; and of a darkish brown colour. This figure of the human gall stones has not ever been rationally accounted for. They vary also much in degree of consistency; some being soft and friable, which

are generally of a light colour.—Some will swim in water; and others sink. In size they have been found to weigh even some ounces, and in number actually many hundreds. Some of these gall stones, when held to a slame take fire and will burn like resin: whereas others will not burn, but melt like wax.—The solid dark coloured ones do not dissolve by gentle coction, either in spirits or water: of some, above one half will dissolve in warm water.

A peculiar and not uncommon appearance of gall stones, is a great number of very minute black ragged granules.

Mr. Cline mentioned to me, that in one case he met with the gall bladder distended with a transparent ropey sluid, not in the least colored with bile, and in which the cystic duct was imperforate. In this gall bladder several small irregular black calculi were contained. This sluid was probably secreted by the internal coat of the gall bladder, which from its great vascularity seems capable of performing that

that function: and it is probable that these calculi were formed by that secretion, consequently not biliary calculi.— Also in another case the cystic duct was entirely obstructed by a calculus, and the gall bladder was distended by the same kind of sluid as the former, and contained the same irregular substances.—In both these cases, bile was contained in the pori biliarii and the biliary ducts, in the usual quantity.—From which circumstances it appears probable that these kind of concretions, are formed by the sluid secreted in the gall bladder.

Biliary calculi are no less various in fize and number, than common as a disease, or appearance after death. Many extraordinary instances of which may be seen* in papers published by the medical societies of most countries.

The

In the works of Galen, Du Verney, Vaterius, Fernelius, Vessingius, Gentili, Nicolus, Fallopius, Haller, Morgagni, Hossman, &c.—Vesalius sound in the body of Prosper Martelli, the gall bladder twice the size of a sist, and filled with calculus granules like the seeds of millet.

The symptoms caused by them vary, from the most violent lancinating and spasmodic pains in the region of the gall bladder, even to the least imaginable inconvenience.—And on the contrary, there is no doubt from the frequent instances of such concretions having been discovered after death, where no bilious affection had preceded, that such may, and do frequently happen without perceptible inconvenience.

Lithotomy has been advised for the removal of these stones. This however has arose from the uncandid representation of a very uncommon case, but which has I know occurred: to wit, of an abscess forming between the calculus, and the integuments of the abdomen. In this

Dr. Hunter in his lectures used to recite the case of a young woman, in whose gall bladder were found 1100 gall stones, and yet no pain or disease of the digestive organs had preceded.

The celebrated Dr. Mead found a gall stone weighing two drams, in the gall bladder of the late Lord Bath, notwithstanding which, neither pain in the part, nor jaundice had preceded,

case

case when the matter is let out, the stone may be felt by a probe or the finger introduced into the wound, and possibly extracted. But the kind and wife process of nature in fuch cases is, first to form a complete adhesion between the gall bladder or ducts and peritonæum, so that when fuch abscess is opened, the cavity of the belly remains unexposed, that is, not cut into. One case of this kind I have feen, in which the abfcess had been opened, and a fiftulous aperture remained leading to a gall stone. In Morgagni de causis and sedibus morborum are three of fuch cases. The first was cured: the fecond had a fiftula left, by which a thin yellow liquid was discharged; the third had an ulcer remain, which with it's fanies, discharged bilious calculi at times.

The remedies for this disorder are laxatives joined with diuretics, neutral salts, emetics, and opiates: solvents have also been recommended. It appears necessary to caution against the indiscriminate use of stimulant purgatives or emetics, from the danger of their encreasing the pain and irritation. For it is to be considered that any stimulating medicine, would by encreasing the irritation of the biliary ducts, tend to exaggerate the symptoms.

Biliary calculi of such remarkably large dimensions have been discharged by stool,* that practitioners were puzzled in endeavouring to account for the course through which they came.—It has been supposed, from the impossibility of their passing through the ducts, that they were generated in the stomach or bowels.—But Mr. Cline having found the gall bladder adhering to the duodenum, and a direct aperture from it into that intestine:— (which parts are among his anatomical preparations:)—It appears that in this

course

^{*} An extraordinary case of this kind is related by Dr. Lettsom, in the first volume of the Memoirs of the Medical Society of London.

course a stone of any dimensions, confined in the gall bladder, might take it's passage directly into the gut, and be discharged by stool.

Opiates properly timed, will be found to be the sovereign remedy in alleviating the pains and spasms occasioned actually by such calculi, or by the obstructed bile itself. But they should be joined with laxatives; and so much of an emetic medicine, as to excite some degree of nausea without vomiting may also be administered with them. The warm bath, or somentations to the hypochondriac region may afford relief.

Solvents for these calculi have likewise been proposed, and are highly extolled by some authors. The juice of common grass has been called a specific; founded on the remark that oxen sed on dry sodder are subject to such calculi; from the symptoms of which seeding on grass cures them.—But when we resect on the course that such medicine must take in the circulation

culation before it arrives at the liver, in order to mix with the bile, and to act as a folvent on these calculi; the difficulty of introducing a fufficient quantity of any fuch medicine will be evident.* Much however may be expected from diet of an attenuant kind, confisting much of vegetables, and the free use of warm diluent drinks are also to be recommended, in order to attenuate the bile; unless there was any evident tendency to dropfy.-In jaundice long continued, and arifing from a viscid state of the bile, an attenuating diet confisting principally of vegetables, may be confidered as the chief means by which a cure is to be expected, for this obvious reason; that the bile being secreted from the blood returned by the veins from the abdomen, it's nature will depend very much upon, and may be very eafily altered by diet. For beverage, hock and water, or in it's stead cyder and water is highly proper, as is water by itself, or a

^{*} A decoction of Pareira Brava is recommended by Hoffman.

light

light infusion of some grateful aromatic or mild bitter; whey is also proper. Sour fruits might also be of service; acids and some of their compounds having the power of decomposing the bile.

On the subject of exercise, the same observations will apply that were made on the use of cathartics and emetics. It is capable of doing harm; as may negatively appear from this circumstance, that bending the body forwards for any length of time, as in writing or reading upon a desk, is particularly injurious in most bilious affections. Gentle exercise on horseback, failing, or fwinging, may be recommended; benefit may also accrue from what is vulgarly called, standing on the head. The criterion by which I should judge of the safety of exercise would be of its encreasing the pain or not. Wherefore, moderate exercise should be first attempted.

^{*} Fordyce Blem. pract. physic.

The next kind of jaundice, is that arifing from the pressure of some indurated part on the biliary ducts. Cases of this kind are to be found in some of the authors already quoted. The fymptoms of this I should suppose are also to be collected negatively, to wit, from the abfence of the fymptoms ordinarily attendant upon jaundice; from it's not being attended with pain in the epigastric, hypochondriae, or umbilical region, (which latter Hoffman afferts to be the most clear indication of gall stones;) from it's not yielding to the remedies recommended for common jaundice, continuing long without intermission, or from it's having been preceded by symptoms of obstructed viscera.-In one case of this kind an incurable jaundice was caused by the presfure of a schirrous pancreas upon the ducts of the gall bladder .- An inflammation happening in the concave part of the liver, is also in some instances productive of jaundice.

The remedies most likely to afford E 2 relief

relief would probably be a mercurial course, cicuta, or the folvens mineralis.

The next cause of jaundice enumerated, is contraction of the biliary ducts themfelves .- One instance of this disease I met with in the examination of the body of a gentleman who died of a fever, but had been previously often afflicted with jaundice.-In this case, the ductus communis cholidochus was near it's opening into the bowel fo contracted, that a large pin would scarcely pass, and such a tenacious fluid as bile with much difficulty. Another proof of this may be drawn from the effect of the mind, in some who are subject to jaundice, in whom anxiety and other passions, will bring on this diforder in a few hours.—Some poisons are also productive of this, which may be called the spasmodic jaundice, as the bites of venomous animals. The bite of the viper produces this kind of jaundice in less than an hour; which can only be accounted for on the theory of every part of the body fecreting bile, in some cases Amilaof jaundice. The principal remedies for this jaundice will be antispasmodics, and even opiates if those should not be successful.

A viscid state of the bile constitutes another cause of jaundice. This when habitual and not amounting to disease, seems to have been described by the ancients under the title of the melancholic temperament. The symptoms are a lassitude of body and dispiritedness, with jaundice of a dark or dingy yellow ‡. This kind of jaundice I should also suffered to be not unfrequently the cause of

^{*} Fordyce Elem. pract. physic.

[†] These ducts have also been found much dilated, even so as to admit a moderate sized singer at the insertion of the ductus communis, into the duodenum.

In the recent jaundice the colour is of a bright light yellow: as happened once to myself, being compleatly jaundiced in one night. This was preceded by a constant pain for several days at the region of the gall bladder, which was relieved from the instant the jaundice took place: owing probably to the absorbents having removed the quantity of bile, which by distending the gall bladder caused the pain.

afthmatic affections. When the species of jaundice can be clearly ascertained, it would be right to make a rather free use of cathartics*, affished by such other medicines as the symptoms may require; especially antispasmodics with opiates; an attenuating diet and the free use of diluting liquids, in order to thin the blood and juices secreted from the same, with moderate exercise, and ease of mind.—Haller says the bile becomes viscid, "quieto, obeso, & a mærore."

Another cause of jaundice, reckoned by some as the most common, is an obstruction of the small glands or biliary follicles in the liver. This however I imagine to be very seldom the real cause of jaundice, which may be more rationally deduced from a viscidity of the bile itself, slowing too slowly through it's natural ducts, and in consequence of this remora, admitting of some absorption of

^{*} Aloes are peculiarly adapted to this diforder, being indued with the property of rarefying the blood. Cullen's Lectures on the Materia Medica.

bile into the blood, and thus inducing jaundice.

ly eartist, as to caude not only feeling

The last general head under which I have classed bilious diseases, is when they are excited by the diseased state of the bile. Such cases I think are not frequent in England, or any of the northern climates.—An acrid state of the bile is capable of stimulating the bowels and stomach, thereby producing sudden and violent evacuations by vomit and stool, then called cholera morbus. Bilious diarrhæa's do also sometimes occur from this cause.

These symptoms have been treated on in some of the preceding pages, as arising from an excessive quantity of bile. It is also more than probable, that these two causes are united in some, perhaps most of such cases.—Fevers from acrimony of bile are also sometimes met with here, particularly in hot and dry seasons.

In hot countries, bile is fometimes accumulated in the intestines, and becomes fo highly putrid, as to cause not only fevers of a most dangerous kind, but various other diseases. But it appears to me, that fuch disorders are much more frequently caused by the contents of the stomach and bowels becoming putrid. For (especially in hot climates) I imagine it may be admitted as a fact in physic, that " animal food will always become putrid, unless it be digested, or else evacuated fooner than is natural," which is to be accounted for, from the stomach in health affimilating the aliment to a pulpy mass, without its going into a state of putrefaction, and having the power of stopping the process of an incipient putrefaction *. -Whereas human bile does not putrify more readily than the other fecretions, or the blood. +

It has been found in India, that the sudden disturbance and evacuation of such

See Spallanzani's experiments on digeftion.

⁺ Fordyce, Elem. physic.

bile, and contents of the primæ viæ, is a matter of great hazard, requiring nice discrimination, as many have died under the evacuations, as supposed from its extreme acrimony. The practitioner's object should therefore be to dilute and correct the ill state of such offending bile and putrid contents of the stomach and bowels, by proper diluents and medicines, previous to its evacuation.

Another general remark.—Chronic bilious diseases, and those of the liver, are preceded by costiveness: are often caused by anxiety of mind, and too frequently by drinking of spirits.—Some of the leading symptoms caused by drinking, are pain in the stomach, loss of appetite, symptoms of asthma, jaundice, and costiveness, the precursors of satal consumption or dropsy.

Vide, Memoirs of the Medical Society of London, vol. 1. page 152. the symptoms arising from drinking, accurately described by Dr. Lettsom, page 153, are these words, "the liver seems less than natural."

It feems but rational to suppose that many of these diseases may be prevented by avoiding these causes; and by a due attention to the state of the bowels, which should be kept open daily.—By this I mean, that the mind should be relieved by company, change of scene, and so on: but that the bottle should be refrained from, as more particularly liable to bring on disease of the liver at this period; fince it has been proved, that anxiety of mind alone is capable of inducing bilious diseases .- I know a gentleman, who, with a constitutional bad digestion, has drank part of a glass of brandy, and a bottle, often two, of wine daily after dinner, with few intermissions, for forty years, without any apparent ill effect: but he has constantly kept his body open, and this he has done chiefly by the means of hiera picra.—This latter observation merits the particular attention of all those who drink freely, for I am persuaded that many liver diseases, which end fatally, might be thus prevented. In general, some warm laxative will be requifite :

fite; that which will agree best may be discovered by each individual, on trying various kinds: fuch as rhubarb, vinum aloes, or any fuch.-I should be forry to be misunderstood, as being the advocate of the free use of the bottle, being certain that it shortens life; meaning what I have here advanced to apply to those who will not lay afide this baneful practice; for such are still entitled to our aid. Further, it is not the quantity alone which destroys the constitution, but the time when drank: for ftrong liquor taken into the stomach when void of food, will exert its heating and corrugating powers over that organ and the liver with infinitely more effect than when it contains folid food, with which it will become well blended, and of consequence not so deleterious to these parts .- It is also highly worthy of remark, that a certain given quantity of strong liquor drank at one or two stated times of a day, (the only proper times are after dinner, and after supper,) injure much less than if indiscriminately taken; whereas drinking

in the morning kills many hundreds yearly in England.

Laftly, Pain at the pit of the stomach. being one very common attendant upon biliary diseases, I judged that the following observation was founded on a fact of too much consequence to be omitted: viz. that the family remedy for this, is fome spirituous cordial.—Language too poignant cannot be held out against this custom, fince to this, the fatal termination of many of these cases in incurable liver disease, dropsy, &c. may be attributed, positively in some instances, I think myself well founded in afferting, more than to the original distemper. Brandy and water is also a common but a most treacherous, beguiling, poisonous, remedy, obnoxious to the same objections, and to the further one of the propenfity which people have to encrease its strength, in proportion to the progressive weakness of the stomach. - From my own experience I found it was the warm water which afforded the relief: a dish

of warm coffee or tea * was my remedy for two years, during which time I was conftantly at noon afflicted with an urgent pain at the stomach.—But let us suppose that I had daily at noon, on an empty stomach during two years, that is more than seven hundred times, taken some cordial or strong drink: is it not more than probable that an incurable liver disease would have been the consequence? I am certain it would.

In the foregoing pages much stress has been placed upon diet: I shall, therefore, beg leave to conclude in the words of the late ingenious Cullen.

"It were well, therefore, that mankind were aware of the tendency which every kind of diet has to produce effects, either immediately, or after repetition, unfavorable to health. And I cannot be of more fervice than by engaging physicians in a minute study of the subject."

FINIS.

^{*} To those who have any dislike to these liquids, I would recommend a light infusion of ginger, orange peel, or any such warm mild bitter, or lemonade.

To the Faculty and Students of Medicine.

The following BOOKS are printed for J. MURRAY,
No. 32, Fleet-Street.

Dedicated by Permission to His Majesty.

I. A N ENQUIRY into the SMALL POX, MEDICAL and POLITICAL; wherein a successful
method of treating that disease is proposed, the cause
of Pits explained, and the Method of their prevention
pointed out; with an Appendix, representing the present State of the Small Pox,

By ROBERT WALKER, M. D. Fellow of the Royal College of Surgeons, Edinburgh.

Octavo, 6s. in boards.

II. A New Edition, being the 4th, corrected throughout, AN ESSAY on DISEASES incidental to EURO-PEANS in HOT CLIMATES; with the method of preventing their fatal consequences,

By JAMES LIND, M.D. F. R.S. Edinburgh, &c. To which is added an Appendix, concerning Intermittent Fevers, and a fimple and easy Way to render Sea Water fresh, and to prevent a Scarcity of Provisions in long Voyages at Sea. Octavo, 5s. boards.

III. DISSERTATIONS relative to the NATURAL HISTORY of ANIMALS and VEGETABLES. Translated from the Italian of the Abbe Spallanzani. Octavo, 2 Vol. 12s. boatds.

IV. TREATISE on the VENOM of the VIPER; on the AMERICAN POISONS, and on the CHERRY LAUREL; and fome other VEGETABLE POISONS. To which are annexed, Observations on the Primitive Structure of the Animal Body; different Experiments on the Reproduction of the Nerves; and a Description of a New Canal of the Eye. With Ten Descriptive Plates. Translated from the original French of Felix Fontana, Naturalist to his Royal Highness the Grand Duke of Tuscany, and Director of his Cabinet of Natural History,

By JOSEPH SKINNER, Navy Surgeon, and Member of the Corporation of Surgeons of London, 2 Vol. Octavo, 125. boards.

V. LETTERS and ESSAYS on the Small Pox and Inoculation, the Measles, the Dry Belly Ache, the Yellow, and Remitting and Intermitting Fevers of the WestWest-Indies. To which are added, Thoughts on the Hydrocephalus, Internus, and Observations on Hydatides in the Heads of Cattle, by different Practitioners. Octavo, price 5s. bound.

VI.MEDICAL COMMENTARIES. Exhibiting a concise view of the latest and most important Discoveries in Medicine and Medical Philosophy, collected and published by ANDREW DUNCAN, M. D. F. R. and A. S. Edin. &c. &c. Octavo, 14 Vol. 41. 18s. bound.

N. B. Any of the Volumes may be had separate.

VII. The DOMESTIC PHYSICIAN; or GUARDIAN of HEALTH; pointing out, in the most familiar manner, the Nature and Symptoms of every Disorder incident to Mankind, with their gradual Progress, and the Method of Cure; particularly adapted to the use of Families and private Practitioners,

By B. CORNWELL, M.L.

*In this Work both the Acute and Chronic Diseases are treated in a clear and comprehensive Manner; it also contains every necessary advice to the Fair Sex, with competent Instructions for the Nursing and Management of Children; Directions for the Cure of Wounds, Fractures, Dislocations, &c. Rules for the Preservation of Health, suited to the various Occupations and Situations of Mankind; with an Appendix forming a complete Dispensary, with a copious Index, Octavo, 7s. 6d. bound.

VIII.EXPERIMENTS & OBSERVATIONS on quilled and red PERUVIAN BARK; among which are included fome remarkable effects arifing from the action of common Bark and Magnesia upon each other; with remarks on the nature and mode of treatment of Fevers, putrid Sore Throat, Rheumatism, Scrophula, and other diseases; in order to ascertain the cases in which Bark may be administered—either alone, or combined with other remedies, to the best advantage. To which is added, an Appendix on the Cinchona Caribboa,

By THOMAS SKEETE, M.D. Octavo, price 5s. in boards.

IX. FORSTER'S PRINCIPLES and PRACTICE of MIDWIFERY, in which are comprised and methodically arranged, under the four general heads of Generation, Gestation, Delivery, and Recovery, all the anatomical facts, physiological reasonings, pathological observations, and practical precepts, necessary to constitute the fullest and

and most complete System of Midwifery. Octavo, price 4s. in boards.

X. HOME's Clinical Experiments, Histories, and Diffections. Second edition. Price 6s. boards.

XI. The CHEMICAL ESSAYS of Charles William Scheele. Translated from the Transactions of the Academy of Sciences at Stockholm, revised, and with additions, by THOMAS BEDDOES, Esq. President of the Royal Medical, of the Natural History, and Chemical Societies at Edinburgh. Octavo, price 6s. in boards.

XII. BERGMAN's Chemical Effays, 2 vols. Translated by Dr. E. Cullen. With Notes. Octavo, price 13s. in boards.

XIII. BERGMAN on the USEFULNESS of CHE-MISTRY. Octavo, 2s. 6d. fewed.

XIV. Brunonis Elementa Medicinae, editio altera pluzimum emendata, Octavo, 2 Vol. 8s. boards.

XV. BROWN'S OBSERVATIONS on the OLD SYSTEM of PHYSIC, exhibiting a Compendium of the New Doctrine. Octavo, 6s. boards.

XVII. MURRAY'S CATALOGUE of BOOKS in MEDICINE, SURGERY, ANATOMY, NATURAL HISTORY, &c. which are now felling (for ready money) at the prices marked against the several articles, by J. Murray, No. 32, Fleet-Street.

